

BENCHMARKS FOR DEVELOPING ECOLOGICAL SOIL SCREENING LEVELS (ECO-SSL): EFFECTS OF SELENIUM ON SOIL INVERTEBRATES

Ronald T. Checkai, Michael Simini, Roman Kuperman,
Carlton T. Phillips, Dennis W. Johnson
U. S. Army Edgewood Chemical Biological Center
Aberdeen Proving Ground, MD 21010-5424 USA

Richard M. Higashi, University of California, Davis, CA 95616 USA
Teresa W.-M. Fan, University of Louisville, Louisville, KY 40208 USA

Keith Sappington, U.S. EPA ORD National Center for Environmental Assessment
Washington DC 20460 USA

ABSTRACT

Army concepts for the Objective Force include the need to be responsive and deployable. Both of these rely upon proper testing and training. Neither testing nor training can occur without environmental compliance with NEPA, that routinely calls for Environmental Assessments including Human Health Risk Assessment and Ecological Risk Assessment. The U.S. Environmental Protection Agency in a collaborative effort of a multi-stakeholder workgroup including Army, Navy, and Air Force, has established national guidance and SOPs for deriving Ecological Soil Screening Levels (Eco-SSL) for ecological receptors. Unfortunately data in the published literature were insufficient in quantity and quality to establish an Eco-SSL for selenium, critical in various DoD advanced technologies, and frequently found in the environment at testing and training areas. The EC₂₀ ecotoxicological benchmarks based on toxicity of selenium to reproduction by the common soil invertebrates earthworm *Eisenia fetida*, potworm *Enchytraeus crypticus*, and collembolan *Folsomia candida*, were remarkably similar yielding the values 3.4, 4.4, and 4.7 mg kg⁻¹, respectively. Selenium benchmarks will be submitted to the Eco-SSL National Task Group for use in developing a selenium Eco-SSL for soil invertebrates. Establishing Eco-SSL leads to increased Army/DoD readiness by supporting environmental documentation required for testing and training.

1. INTRODUCTION

The United States Army has delineated concepts for the Objective Force, and among these are the need to be responsive and deployable. Both responsiveness and deployability rely upon proper testing and training. Neither testing nor training can occur without

environmental compliance with the National Environmental Policy Act (NEPA). The Army complies with NEPA requirements as directed in 32 CFR 651 / 65 FR 54348. NEPA routinely calls for Environmental Assessments (EA), including Human Health Risk Assessment and Ecological Risk Assessment (ERA). The U.S. Environmental Protection Agency (USEPA) in a collaborative effort of a multi-stakeholder workgroup consisting of USEPA, DoD (Army, Navy, and Air Force), Department of Energy, states, universities, and consulting and industry representatives, has established national guidance and SOPs (USEPA, 2003) for deriving Ecological Soil Screening Levels (Eco-SSL) for ecological receptors including soil invertebrates. Eco-SSLs are screening ecotoxicity values that can be used routinely to identify contaminants of potential concern in soils that warrant additional evaluation in Baseline ERA, and to eliminate those that do not.

Unfortunately data in the published literature were insufficient in quantity and quality to establish Eco-SSL for all of the most common contaminants of ecological and military importance. The metal selenium is a critical element in various Department of Defense (DoD) advanced technologies, and is frequently found in the environment at DoD testing and training areas. These investigations were undertaken to establish ecotoxicological benchmarks that meet Eco-SSL criteria for deriving screening levels based on the toxicity of selenium to soil invertebrates.

2. EXPERIMENTAL RESULTS AND DISCUSSION

We determined EC₂₀ ecotoxicological benchmarks based on reproduction (juveniles produced) for the toxicity of selenium to the soil invertebrates earthworm *Eisenia fetida*, potworm *Enchytraeus crypticus*, and collembolan *Folsomia candida*. Sassafras sandy loam [Fine-loamy,

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 00 DEC 2004		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Benchmarks For Developing Ecological Soil Screening Levels (Eco-Ssl): Effects Of Selenium On Soil Invertebrates				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U. S. Army Edgewood Chemical Biological Center Aberdeen Proving Ground, MD 21010-5424 USA; University of California, Davis, CA 95616 USA				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001736, Proceedings for the Army Science Conference (24th) Held on 29 November - 2 December 2005 in Orlando, Florida. , The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

siliceous, mesic Typic Hapludult] at pH 7.1, with sand:silt:clay ratio 70:13:17, and 1% organic matter, was utilized in the bioassays. Selenium was added as sodium selenate, and weathered/aged in soil via alternating wetting/drying cycles for 13 weeks. Soil from each treatment level was collected, lyophilized, and stored at -40°C at the initiation and conclusion of each test. Soils were analyzed for selenium by LC/ICP-MS, following nitric:perchloric digestion. Nonlinear regression models were used to correlate ecotoxicity data with selenium concentrations, retained primarily as SeO_4 in this aerobic soil. The benchmarks for selenium were remarkably similar, yielding EC_{20} values 3.4, 4.4, and 4.7 mg kg^{-1} for earthworm, potworm, and collembola, respectively. Selenium benchmarks will be submitted to the Eco-SSL National Task Group for use in developing an Eco-SSL for soil invertebrates; experimental design and results undergo quality assurance before inclusion of benchmarks in the national Eco-SSL database.

3. CONCLUSIONS

Eco-SSL eliminate the need for repetitious costly toxicity-data literature searches; increase consistency; and allow resources to be focused on issues critical to sustaining key testing and training sites. Establishing Eco-SSL leads to increased Army/DoD efficiency and readiness by supporting environmental documentation required for testing and training, essential for force responsiveness and deployability.

ACKNOWLEDGEMENT

This research supported by U.S. Environmental Protection Agency, Interagency Agreement No. DW-21-93926501-0.

REFERENCE

USEPA, 2003: Ecological Soil Screening Level Guidance. U.S. Environmental Protection Agency, <http://www.epa.gov/ecotox/ecossl/>, Washington, DC.